

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

**EP 0 906 959 A2**

(12)

**EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
07.04.1999 Bulletin 1999/14

(21) Application number: 98305919.7

(22) Date of filing: 24.07.1998

(51) Int Cl.<sup>6</sup>: **C12N 15/57**, C12N 9/48,  
C07K 16/40, C12P 21/02,  
C12N 5/10, C12Q 1/37,  
A61K 38/48, A61K 39/085,  
A61K 48/00

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU**  
**MC NL PT SE**  
Designated Extension States:  
**AL LT LV MK RO SI**

(30) Priority: 05.08.1997 US 55388 P  
05.11.1997 US 964631

(71) Applicants:  
• **SMITHKLINE BEECHAM CORPORATION**  
Philadelphia Pennsylvania 19103 (US)  
• **SMITHKLINE BEECHAM PLC**  
Brentford, Middlesex TW8 9EP (GB)

(72) Inventors:  
• **Lonetto, Michael Arthur**  
King of Prussia, Pennsylvania 19406 (US)  
• **Warren, Patrick Vernon**  
King of Prussia, Pennsylvania 19406 (US)  
• **Biswas Sanjoy**  
King of Prussia, Pennsylvania 19406 (US)  
• **Warren, Richard Lloyd**  
King of Prussia, Pennsylvania 19406 (US)  
• **Burnham, Martin K.R.**  
Brentford, Middlesex, TW8 9EP (GB)

(74) Representative: **Mallalieu, Catherine Louise et al**  
**D. Young & Co.,**  
21 New Fetter Lane  
London EC4A 1DA (GB)

(54) **Aminopeptidase from Staphylococcus aureus**

(57) The invention provides aminopeptidase polypeptides and DNA (RNA) encoding aminopeptidase polypeptides and methods for producing such polypep-

tides by recombinant techniques. Also provided are methods for utilizing aminopeptidase polypeptides to screen for antibacterial compounds.

**EP 0 906 959 A2**

**Description****RELATED APPLICATIONS**

5 [0001] This application claims benefit of US Provisional Patent application number 60/055,388, filed August 5, 1997

**FIELD OF THE INVENTION**

10 [0002] This invention relates to newly identified polynucleotides and polypeptides, and their production and uses, as well as their variants, agonists and antagonists, and their uses. In particular, in these and in other regards, the invention relates to novel polynucleotides and polypeptides of the M29 aminopeptidase family, hereinafter referred to as "aminopeptidase".

**BACKGROUND OF THE INVENTION**

15 [0003] It is particularly preferred to employ Staphylococcal genes and gene products as targets for the development of antibiotics. The Staphylococci make up a medically important genera of microbes. They are known to produce two types of disease, invasive and toxigenic. Invasive infections are characterized generally by abscess formation effecting both skin surfaces and deep tissues. *S. aureus* is the second leading cause of bacteremia in cancer patients. Osteomyelitis, septic arthritis, septic thrombophlebitis and acute bacterial endocarditis are also relatively common. There are at least three clinical conditions resulting from the toxigenic properties of Staphylococci. The manifestation of these diseases result from the actions of exotoxins as opposed to tissue invasion and bacteremia. These conditions include: Staphylococcal food poisoning, scalded skin syndrome and toxic shock syndrome

20 [0004] The frequency of *Staphylococcus aureus* infections has risen dramatically in the past 20 years. This has been attributed to the emergence of multiply antibiotic resistant strains and an increasing population of people with weakened immune systems. It is no longer uncommon to isolate *Staphylococcus aureus* strains which are resistant to some or all of the standard antibiotics. This has created a demand for both new anti-microbial agents and diagnostic tests for this organism.

25 [0005] AMPS proteins are members of the M29 peptidase family. These are broad spectrum aminopeptidases using cobalt and magnesium as cofactors. These proteins represent possible targets for novel antibacterials as their inhibition would interrupt the tissue damaging actions of pathogens, leading to their inability to promote and maintain infection.

30 [0006] Substantial effort has been invested this century in the successful discovery and development of antibacterials. Paradoxically although antibacterials are devised to eradicate infection in mammals we know almost nothing of the physiology of bacterial pathogens in infective situations in the host. Using sequences from the *Staphylococcus aureus* chromosome we have developed an RT-PCR based procedure which allows us to identify those bacterial genes transcribed at any stage of infection and also from different niches of infection. The derivation of such information is a critical first step in understanding the global response of the bacterial gene complement to the host environment. From the knowledge of bacterial genes both of known and unknown function which are widely transcribed in the host it is possible to attempt to ascertain by database searching those which are present only in the eubacteria. Further prioritisation of such genes by consideration of the likely role of their products towards the maintenance of infection and the facility of setting up a screen for inhibitors of the biochemical function indicated by their homology to characterised genes allows the compilation of a shortlist for gene essentiality studies using genetic deletion or controlled regulation techniques. The proteins expressed by genes shown to be necessary for growth *in vitro* or in pathogenesis in animal models provide novel targets for antibacterial screening to find agents which are broadly inhibitory towards pathogenesis. This invention provides *S. aureus* WCUH 29 polynucleotides which are transcribed in infected tissue, in particular in both acute and chronic infections.

40 [0007] Clearly, there is a need for factors, such as the novel compounds of the invention, that have a present benefit of being useful to screen compounds for antibiotic activity. Such factors are also useful to determine their role in pathogenesis of infection, dysfunction and disease. There is also a need for identification and characterization of such factors and their antagonists and agonists which can play a role in preventing, ameliorating or correcting infections, dysfunctions or diseases.

50 [0008] The polypeptides of the invention have amino acid sequence homology to a known AMPS\_BACSU (SWISS-PROT: locus AMPS\_BACSU, accession P39762) protein

**SUMMARY OF THE INVENTION**

55 [0009] It is an object of the invention to provide polypeptides that have been identified as novel aminopeptidase polypeptides by homology between the amino acid sequence set out in Table 1 [SEQ ID NO: 2] and a known amino

acid sequence or sequences of other proteins such as AMPS\_BACSU (SWISS-PROT: locus AMPS\_BACSU, accession P39762) protein.

[0010] It is a further object of the invention to provide polynucleotides that encode aminopeptidase polypeptides, particularly polynucleotides that encode the polypeptide herein designated aminopeptidase.

5 [0011] In a particularly preferred embodiment of the invention the polynucleotide comprises a region encoding aminopeptidase polypeptides comprising the sequence set out in Table 1 [SEQ ID NO: 1] which includes a full length gene, or a variant thereof.

[0012] In another particularly preferred embodiment of the invention there is a novel aminopeptidase protein from *Staphylococcus aureus* comprising the amino acid sequence of Table I [SEQ ID NO:2], or a variant thereof.

10 [0013] In accordance with another aspect of the invention there is provided an isolated nucleic acid molecule encoding a mature polypeptide expressible by the *Staphylococcus aureus* WCUH 29 strain contained in the deposited strain.

[0014] A further aspect of the invention there are provided isolated nucleic acid molecules encoding aminopeptidase, particularly *Staphylococcus aureus* aminopeptidase, including mRNAs, cDNAs, genomic DNAs. Further embodiments of the invention include biologically, diagnostically, prophylactically, clinically or therapeutically useful variants thereof, and compositions comprising the same.

[0015] In accordance with another aspect of the invention, there is provided the use of a polynucleotide of the invention for therapeutic or prophylactic purposes, in particular genetic immunization. Among the particularly preferred embodiments of the invention are naturally occurring allelic variants of aminopeptidase and polypeptides encoded thereby.

20 [0016] Another aspect of the invention there are provided novel polypeptides of *Staphylococcus aureus* referred to herein as aminopeptidase as well as biologically, diagnostically, prophylactically, clinically or therapeutically useful variants thereof, and compositions comprising the same.

[0017] Among the particularly preferred embodiments of the invention are variants of aminopeptidase polypeptide encoded by naturally occurring alleles of the aminopeptidase gene.

25 [0018] In a preferred embodiment of the invention there are provided methods for producing the aforementioned aminopeptidase polypeptides.

[0019] In accordance with yet another aspect of the invention, there are provided inhibitors to such polypeptides, useful as antibacterial agents, including, for example, antibodies.

30 [0020] In accordance with certain preferred embodiments of the invention, there are provided products, compositions and methods for assessing aminopeptidase expression, treating disease, for example, disease, such as, infections of the upper respiratory tract (e.g., otitis media, bacterial tracheitis, acute epiglottitis, thyroiditis), lower respiratory (e.g., empyema, lung abscess), cardiac (e.g., infective endocarditis), gastrointestinal (e.g., secretory diarrhoea, splenic abscesses, retroperitoneal abscess), CNS (e.g., cerebral abscess), eye (e.g., blepharitis, conjunctivitis, keratitis, endophthalmitis, preseptal and orbital cellulitis, dacryocystitis), kidney and urinary tract (e.g., epididymitis, intrarenal and perinephric abscess, toxic shock syndrome), skin (e.g., impetigo, folliculitis, cutaneous abscesses, cellulitis, wound infection, bacterial myositis) bone and joint (e.g., septic arthritis, osteomyelitis), assaying genetic variation, and administering a aminopeptidase polypeptide or polynucleotide to an organism to raise an immunological response against a bacteria, especially a *Staphylococcus aureus* bacteria.

40 [0021] In accordance with certain preferred embodiments of this and other aspects of the invention there are provided polynucleotides that hybridize to aminopeptidase polynucleotide sequences, particularly under stringent conditions.

[0022] In certain preferred embodiments of the invention there are provided antibodies against aminopeptidase polypeptides.

45 [0023] In other embodiments of the invention there are provided methods for identifying compounds which bind to or otherwise interact with and inhibit or activate an activity of a polypeptide or polynucleotide of the invention comprising: contacting a polypeptide or polynucleotide of the invention with a compound to be screened under conditions to permit binding to or other interaction between the compound and the polypeptide or polynucleotide to assess the binding to or other interaction with the compound, such binding or interaction being associated with a second component capable of providing a detectable signal in response to the binding or interaction of the polypeptide or polynucleotide with the compound; and determining whether the compound binds to or otherwise interacts with and activates or inhibits an activity of the polypeptide or polynucleotide by detecting the presence or absence of a signal generated from the binding or interaction of the compound with the polypeptide or polynucleotide.

[0024] In accordance with yet another aspect of the invention, there are provided aminopeptidase agonists and antagonists, preferably bacteriostatic or bacteriocidal agonists and antagonists.

55 [0025] In a further aspect of the invention there are provided compositions comprising a aminopeptidase polynucleotide or a aminopeptidase polypeptide for administration to a cell or to a multicellular organism.

[0026] Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following descriptions and from reading the other parts of the

present disclosure.

## GLOSSARY

[0027] The following definitions are provided to facilitate understanding of certain terms used frequently herein.

[0028] "Host cell" is a cell which has been transformed or transfected, or is capable of transformation or transfection by an exogenous polynucleotide sequence.

[0029] "Identity," as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. "Identity" and "similarity" can be readily calculated by known methods, including but not limited to those described in (*Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data*, Part I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., SIAM J. Applied Math., 48: 1073 (1988). Preferred methods to determine identity are designed to give the largest match between the sequences tested. Methods to determine identity and similarity are codified in publicly available computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, the GCG program package (Devereux, J., et al., *Nucleic Acids Research* 12(1): 387(1984)), BLASTP, BLASTN, and FASTA (Atschul, S.F. et al., *J. Molec. Biol.* 215: 403-410 (1990)). The BLAST X program is publicly available from NCBI and other sources (*BLAST Manual*, Altschul, S., et al., NCBI NLM NIH Bethesda, MD 20894; Altschul, S., et al., *J. Mol. Biol.* 215: 403-410 (1990)). As an illustration, by a polynucleotide having a nucleotide sequence having at least, for example, 95% "identity" to a reference nucleotide sequence of SEQ ID NO: 1 it is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence of SEQ ID NO: 1. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5 or 3 terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence. Analogously, by a polypeptide having an amino acid sequence having at least, for example, 95% identity to a reference amino acid sequence of SEQ ID NO:2 is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of SEQ ID NO: 2. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

[0030] "Isolated" means altered "by the hand of man" from its natural state, i.e., if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living organism is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein.

[0031] "Polynucleotide(s)" generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotide(s)" include, without limitation, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions or single-, double- and triple-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded, or triple-stranded regions, or a mixture of single- and double-stranded regions. In addition, "polynucleotide" as used herein refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The strands in such regions may be from the same molecule or from different molecules. The regions may include all of one or more of the molecules, but more typically involve only a region of some of the molecules. One of the molecules of a triple-helical region often is an oligonucleotide. As used herein, the term "polynucleotide(s)" also includes DNAs or RNAs as described above that contain one or more modified bases. Thus, DNAs or RNAs with backbones modified for stability or for other reasons are "polynucleotide(s)" as that term is intended herein. Moreover, DNAs or RNAs comprising unusual bases, such as

inosine, or modified bases, such as tritylated bases, to name just two examples, are polynucleotides as the term is used herein. It will be appreciated that a great variety of modifications have been made to DNA and RNA that serve many useful purposes known to those of skill in the art. The term "polynucleotide(s)" as it is employed herein embraces such chemically, enzymatically or metabolically modified forms of polynucleotides, as well as the chemical forms of DNA and RNA characteristic of viruses and cells, including, for example, simple and complex cells. "Polynucleotide(s)" also embraces short polynucleotides often referred to as oligonucleotide(s).

[0032] "Polypeptide(s)" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds. "Polypeptide(s)" refers to both short chains, commonly referred to as peptides, oligopeptides and oligomers and to longer chains generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene encoded amino acids. "Polypeptide(s)" include those modified either by natural processes, such as processing and other post-translational modifications, but also by chemical modification techniques. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature, and they are well known to those of skill in the art. It will be appreciated that the same type of modification may be present in the same or varying degree at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains, and the amino or carboxyl termini. Modifications include, for example, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins, such as arginylation, and ubiquitination. See, for instance, *PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES*, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993) and Wold, F., *Posttranslational Protein Modifications: Perspectives and Prospects*, pgs. 1-12 in *POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS*, B. C. Johnson, Ed., Academic Press, New York (1983); Seifter et al., *Meth. Enzymol.* 182:626-646 (1990) and Rattan et al., *Protein Synthesis: Posttranslational Modifications and Aging*, Ann. N.Y. Acad. Sci. 663: 48-62 (1992). Polypeptides may be branched or cyclic, with or without branching. Cyclic, branched and branched circular polypeptides may result from post-translational natural processes and may be made by entirely synthetic methods, as well.

[0033] "Variant(s)" as the term is used herein, is a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques, by direct synthesis, and by other recombinant methods known to skilled artisans.

## DESCRIPTION OF THE INVENTION

[0034] The invention relates to novel aminopeptidase polypeptides and polynucleotides as described in greater detail below. In particular, the invention relates to polypeptides and polynucleotides of a novel aminopeptidase of *Staphylococcus aureus*, which is related by amino acid sequence homology to AMPS\_BACSU (SWISS-PROT: locus AMPS\_BACSU, accession P39762) polypeptide. The invention relates especially to aminopeptidase having the nucleotide and amino acid sequences set out in Table 1 [SEQ ID NO: 1] and Table 1 [SEQ ID NO: 2] respectively, and to the aminopeptidase nucleotide sequences of the DNA in the deposited strain and amino acid sequences encoded thereby.

TABLE 1

## aminopeptidase Polynucleotide and Polypeptide Sequences

(A) Sequences from *Staphylococcus aureus* aminopeptidase polynucleotide sequence [SEQ ID NO:1].

5' -

GTGGGAATTTTGACTAATTACAAAGAAAAGTTACAACAATACGCTGAACTATTAGTTAAGGTTGGTATGA  
ATGTTCAACCAAAACAGCCAGTATTTATTAGATCATCAGTAGAAATATTAGAATTAACGCATTTAATTGT  
AGAAGAAGCATATCATTGTGGCGCTTCAGATGTACGGGTTGTATATAGCGATCCAACTCTTAAAACGCTT  
AAATTTGAAAATGAATCTGTGGAACATTTTGCTAATCATGAGATAAAATCATATGATGTTGAAGCGAGAA  
TGGATTACGTTAAACGTGGCGCTGCTAATTTAGCACTTATTAGTGAAGATCCTGATTTAATGGATAGTAT  
CGATAGCCAAAAATTACAAGCATTTCAACAACAAAACGCACGCGCATTTAAAGGATATATGGAAAGTGTA  
CAAAAAAATCAATTTCCATGGGTCGTAGCTGCTTTTCCATCTAAAGCTTGGGCGAAGAGAGTATATCCTG  
AACTATCTGTAGAAGAAGCATATATTAAGTTTATTGACGAGGTATTTGATATTGTGCGAATTGATGGTAA

CGATCCAGTTGAAAATTGGCGTCAACACATCGCTAATTTAAGTGTTTACGCTCAATATTTACAACAAAA  
AATTACCATGCGTTGCATGATGTATCTGAAGGTACTGATTTAACAGTATGTTTAGCGAAAAATCATATCT  
GGGAAGATGCTACAAGTTATGTTAACGGCGAGGAACAAGCTTTTATTGCTAATATACCAACTGAAGAAGT  
ATTTACAGCACCAGATCGAAATAGAGTACATGGATATGTCACATAATAAATTGCCTTTAAGTTATAATGGA  
ACGATTATCGATCAATTTAAATTGATGTTTAAAGATGGAGAAATTATTGATTTTTCAGCTGAAAAAGGTG  
AAGCGGTATTGAAAGATTTAATCAATACCGATGAAGGTTCAAGAAGATTAGGTGAAGTAGCGTTAGTACC  
TGATGATTCACCGATATCAAATCGTAATACCATTTTTTACAATACTTTATTTGATGAAAATGCAGCATGT  
CATTTAGCGATTGGATCTGCTTACGCCTTTAATATTCAAGGTGGAACGGAAATGACTGTTGAGGAAAAGA  
TTGCAAGCGGATTAAATGATTCAAACGTACATGTCGATTTTCATGATTGGTAGTAGTATTTGACTATTTA  
TGGCATATTCGAAGATGGTTCAAAAGAACTAGTATTTGAAAATGGAAATTGGGCATCAACATTTTAA-3'

(B) aminopeptidase polypeptide sequence deduced from the polynucleotide sequence in this table [SEQ ID NO:2].

NH<sub>2</sub>-

VGILTNYKEKLQQYAELLVKVGMNVQPKQPVFIRSSVEILELTHLIVEEAYHCGASDVRVVYSDPTLKTL  
KFENESVEHFANHEIKSYDVEARMDYVKRGAANLALISEDPLMDSIDSQKLQAFQQNARAFKGYMESV  
QKNQFPWVVAFFPSKAWAKRVPELSVEEAYIKFIDEVFDIVRIDGNDPVENWRQHIANLSVYAQYLQOK  
NYHALHDVSEGTDLTVCCLAKNHIWEDATSYVNGEEQAFIANIPTEEVFTAPDRNRVHGYVTNKLPLSYNG  
TIIDQFKLMFKDGEIIDFSAEKGEAVLKDLINTDEGSRRLEGEVALVPDDSPISNRNTIFYNTLFDEN AAC  
HLAIGSAYAFNIQGGTEMTVEEKIASGLNDSNVHVD FMIGSSDLTIYGI FEDGSKELVFENGWASTF-  
COOH

(C) Polynucleotide sequence embodiments [SEQ ID NO:1].

X-(R<sub>1</sub>)<sub>n</sub>-

GTGGGAATTTTGACTAATTACAAAGAAAAGTTACAACAATACGCTGAACTATTAGTTAAGGTTGGTATGA  
ATGTTCAACCAAAACAGCCAGTATTTATTAGATCATCAGTAGAAATATTAGAATTAACGCATTTAATTGT  
AGAAGAAGCATATCATTGTGGCGCTTCAGATGTACGGGTTGTATATAGCGATCCAACCTCTTAAAACGCTT  
AAATTTGAAAATGAATCTGTGGAACATTTTGCTAATCATGAGATAAAATCATATGATGTTGAAGCGAGAA  
TGGATTACGTTAAACGTGGCGCTGCTAATTTAGCACTTATTAGTGAAGATCCTGATTTAATGGATAGTAT  
CGATAGCCAAAAATTACAAGCATTTCAACAACAAAACGCACGCGCATTTAAAGGATATATGGAAAGTGTA  
CAAAAAAATCAATTTCCATGGGTCGTAGCTGCTTTTCCATCTAAAGCTTGGGCGAAGAGAGTATATCCTG  
AACTATCTGTAGAAGAAGCATATATTAAGTTTATTGACGAGGTATTTGATATTGTGCGAATTGATGGTAA  
CGATCCAGTTGAAAATTGGCGTCAACACATCGCTAATTTAAGTGTTTACGCTCAATATTTACAACAAAAA  
AATTACCATGCGTTGCATGATGTATCTGAAGGTACTGATTTAACAGTATGTTTAGCGAAAAATCATATCT  
GGGAAGATGCTACAAGTTATGTTAACGGCGAGGAACAAGCTTTTATTGCTAATATACCAACTGAAGAAGT  
ATTTACAGCACCAGATCGAAATAGAGTACATGGATATGTCACTAATAAATTGCCTTTAAGTTATAATGGA  
ACGATTATCGATCAATTTAAATTGATGTTTAAAGATGGAGAAATTATTGATTTTTTACGCTGAAAAAGGTG  
AAGCGGTATTGAAAGATTTAATCAATACCGATGAAGGTTCAAGAAGATTAGGTGAAGTAGCGTTAGTACC  
TGATGATTCACCGATATCAAATCGTAATACCATTTTTTACAATACTTTATTTGATGAAAATGCAGCATGT  
CATTTAGCGATTGGATCTGCTTACGCCCTTTAATATTCAAGGTGGAACGGAAATGACTGTTGAGGAAAAGA  
TTGCAAGCGGATTAAATGATTCAAACGTACATGTCGATTCATGATTGGTAGTAGTGAATTTGACTATTTA  
TGGCATATTGGAAGATGGTTCPAAPGAAGTAGTATTTGAAAATGGAAATTGGGCATCAACATTTTAA-

(R<sub>2</sub>)<sub>n</sub>-Y

## (D) Polypeptide sequence embodiments [SEQ ID NO:2].

X-(R<sub>1</sub>)<sub>n</sub>-

5 VGILTNYKEKLQQYAELLVKVGMNVQPKQPVFIRSSVEILELTHLIVEEAYHCGASDVRVVS DPTLKT  
 KFENESVEHFANHEIKSYDVEARMDYVKRGAANLALISEDPLMDSIDSQKLQAFQQQNARAFKGYMESV  
 10 QKNQFPWVVAAFPSKAWAKRVPELSVEEAYIKFIDEVFDIVRIDGNPVENWRQHIANLSVYAQYLQOK  
 NYHALHDVSEGTDLTVCLAKNHIWEDATSYVNGEEQAFIANIPTEEVFTAPDRNRVHGYVTNKLPLSYNG  
 TIIDQFKLMFKDGEIIDFSAEKGEAVLKDLINTDEGSRRLGEVALVPDDSPISNRNTIFYNTLFDEN AAC  
 HLAIGSAYAFNIQGGTEMTVEEKIASGLNDSNVHVFDMIGSSDLTIYGIFEDGSKELVFENGWASTF-  
 (R<sub>2</sub>)<sub>n</sub>-Y

## (E) Sequences from Staphylococcus aureus aminopeptidase polynucleotide ORF sequence [SEQ ID NO:3].

20 5'-CTAATTANCA AAGAAAAGTT ACAACAATAC GCTGAACTAT TAGTTAAGGT TGGTATGAAT  
 GTTCAACCAA AACAGCCAGT ATTTATTAGA TCATCAGTAG AAACATTAGA ATTAACGCAT  
 TTAATTGTAG AAGAAGCATA TCATTGTGGC GCTTCAGATG TACGTGTTGT ATATAGCGAT  
 CCAACCTTAA AACGCTTAAA ATTTGAAAAT GAATCTGTGG AACATTTTGC TAATCATGAG  
 25 ATAAAATCAT ATGATGTTGA AGCGAGAATG GATTACGTTA AACGTGGCGC TGCTAATTTA  
 GCACTTATTA GTGAAGATCC TGATTTAATG GATGGTATCG ATAGCCAAAA ATTACAAGCA  
 TTTCACAAC AAAACGCACG CGCATTTAAA GGATATATGG AAAGTG TACA AAAAAATCAA  
 TTTCCATGGG TCGTAGCTGC TTTTCCATCT AAAGCTTGGG CGAAGAGAGT ATATCCTGAA  
 30 CTATCTGTAG AAGAAGCATA TATTAAGTTT ATTGACGAGG TATTTGATAT TGTGCGAATT  
 GATGGTAACG ATCCAGTTGA AAATTGGCGT CAACACATCG CTAATTTAAG TGTTTACGCT  
 CAATATTTAC AACAAAAAAA TTACCATGCG TTGCATGATG TATCTGAAGG TACTGATTTA  
 35 ACAGTATGTT TAGCGAAAAA TCATATCTGG GAAGATGCTA CAAGTTATGT TAACGGCGAG  
 GAACAAGCTT TTATTGCTAA TATACCAACT GAAGAAGTAT TTACAGCACC AGATCGAAAT  
 AGAGTACATG GATATGTCAC TAATAAATTG CCTTTAAATT TATTTGGAGC GATATTCGAT  
 CAATTTAAAT TGATGTTC-3'

(F) aminopeptidase polypeptide sequence deduced from the polynucleotide ORF sequence in this table [SEQ ID NO:4].

5 NH<sub>2</sub>-Ile Xaa Lys Glu Lys Leu Gln Gln Tyr Ala Glu Leu Leu Val Lys Val  
 1 5 10 15  
 Gly Met Asn Val Gln Pro Lys Gln Pro Val Phe Ile Arg Ser Ser Val  
 20 25 30  
 10 Glu Thr Leu Glu Leu Thr His Leu Ile Val Glu Glu Ala Tyr His Cys  
 35 40 45  
 Gly Ala Ser Asp Val Arg Val Val Tyr Ser Asp Pro Thr Leu Lys Arg  
 15 50 55 60  
 Leu Lys Phe Glu Asn Glu Ser Val Glu His Phe Ala Asn His Glu Ile  
 65 70 75 80  
 Lys Ser Tyr Asp Val Glu Ala Arg Met Asp Tyr Val Lys Arg Gly Ala  
 20 85 90 95  
 Ala Asn Leu Ala Leu Ile Ser Glu Asp Pro Asp Leu Met Asp Gly Ile  
 100 105 110  
 Asp Ser Gln Lys Leu Gln Ala Phe Gln Gln Gln Asn Ala Arg Ala Phe  
 25 115 120 125  
 Lys Gly Tyr Met Glu Ser Val Gln Lys Asn Gln Phe Pro Trp Val Val  
 130 135 140  
 30 Ala Ala Phe Pro Ser Lys Ala Trp Ala Lys Arg Val Tyr Pro Glu Leu  
 145 150 155 160  
 Ser Val Glu Glu Ala Tyr Ile Lys Phe Ile Asp Glu Val Phe Asp Ile  
 165 170 175  
 35 Val Arg Ile Asp Gly Asn Asp Pro Val Glu Asn Trp Arg Gln His Ile  
 180 185 190  
 Ala Asn Leu Ser Val Tyr Ala Gln Tyr Leu Gln Gln Lys Asn Tyr His  
 195 200 205  
 40 Ala Leu His Asp Val Ser Glu Gly Thr Asp Leu Thr Val Cys Leu Ala  
 210 215 220  
 Lys Asn His Ile Trp Glu Asp Ala Thr Ser Tyr Val Asn Gly Glu Glu  
 225 230 235 240  
 45 Gln Ala Phe Ile Ala Asn Ile Pro Thr Glu Glu Val Phe Thr Ala Pro  
 245 250 255  
 Asp Arg Asn Arg Val His Gly Tyr Val Thr Asn Lys Leu Pro Leu Asn  
 260 265 270  
 50 Leu Phe Gly Ala Ile Phe Asp Gln Phe Lys Leu Met Phe  
 275 280 285

55

-COOH

**Deposited materials**

[0035] A deposit containing a *Staphylococcus aureus* WCUH 29 strain has been deposited with the National Collections of Industrial and Marine Bacteria Ltd. (herein "NCIMB"), 23 St. Machar Drive, Aberdeen AB2 1RY, Scotland on 11 September 1995 and assigned NCIMB Deposit No. 40771, and is referred to as *Staphylococcus aureus* WCUH29 on deposit. The *Staphylococcus aureus* strain deposit is referred to herein as "the deposited strain" or as "the DNA of the deposited strain."

[0036] The deposited strain contains the full length aminopeptidase gene. The sequence of the polynucleotides contained in the deposited strain, as well as the amino acid sequence of the polypeptide encoded thereby, are controlling in the event of any conflict with any description of sequences herein.

[0037] The deposit of the deposited strain has been made under the terms of the Budapest Treaty on the International Recognition of the Deposit of Micro-organisms for Purposes of Patent Procedure. The strain will be irrevocably and without restriction or condition released to the public upon the issuance of a patent. The deposited strain is provided merely as convenience to those of skill in the art and is not an admission that a deposit is required for enablement, such as that required under 35 U.S.C. §112.

[0038] A license may be required to make, use or sell the deposited strain, and compounds derived therefrom, and no such license is hereby granted.

**Polypeptides**

[0039] An initial amino acid encoded by the codon "GTG" is indicated in SEQ ID :NO:2 as valine. However, in one embodiment of the invention, this initial amino acid of a polypeptide of the invention is methionine.

[0040] The polypeptides of the invention include the polypeptide of Table 1 [SEQ ID NO:2] (in particular the mature polypeptide) as well as polypeptides and fragments, particularly those which have the biological activity of aminopeptidase, and also those which have at least 70% identity to a polypeptide of Table 1 [SEQ ID NOS:2 and 4] or the relevant portion, preferably at least 80% identity to a polypeptide of Table 1 [SEQ ID NOS:2 and 4], and more preferably at least 90% similarity (more preferably at least 90% identity) to a polypeptide of Table 1 [SEQ ID NOS:2 and 4] and still more preferably at least 95% similarity (still more preferably at least 95% identity) to a polypeptide of Table 1 [SEQ ID NOS:2 and 4] and also include portions of such polypeptides with such portion of the polypeptide generally containing at least 30 amino acids and more preferably at least 50 amino acids.

[0041] The invention also includes polypeptides of the formula set forth in Table 1(D) [SEQ ID NO:2] wherein, at the amino terminus, X is hydrogen, and at the carboxyl terminus, Y is hydrogen or a metal, R<sub>1</sub> and R<sub>2</sub> is any amino acid residue, and n is an integer between 1 and 1000. Any stretch of amino acid residues denoted by either R group, where R is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer.

[0042] A fragment is a variant polypeptide having an amino acid sequence that entirely is the same as part but not all of the amino acid sequence of the aforementioned polypeptides. As with aminopeptidase polypeptides fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most preferably as a single continuous region, a single larger polypeptide.

[0043] Preferred fragments include, for example, truncation polypeptides having a portion of an amino acid sequence of Table 1 [SEQ ID NOS:2 and 4], or of variants thereof, such as a continuous series of residues that includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus. Degradation forms of the polypeptides of the invention in a host cell, particularly a *Staphylococcus aureus*, are also preferred. Further preferred are fragments characterized by structural or functional attributes such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions.

[0044] Also preferred are biologically active fragments which are those fragments that mediate activities of aminopeptidase, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also included are those fragments that are antigenic or immunogenic in an animal, especially in a human. Particularly preferred are fragments comprising receptors or domains of enzymes that confer a function essential for viability of *Staphylococcus aureus* or the ability to initiate, or maintain cause disease in an individual, particularly a human.

[0045] Variants that are fragments of the polypeptides of the invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, these variants may be employed as intermediates for producing the full-length polypeptides of the invention.

**Polynucleotides**

[0046] Another aspect of the invention relates to isolated polynucleotides, including the full length gene, that encode

the aminopeptidase polypeptide having a deduced amino acid sequence of Table 1 [SEQ ID NOS:2 and 4] and polynucleotides closely related thereto and variants thereof.

[0047] Using the information provided herein, such as a polynucleotide sequence set out in Table 1 [SEQ ID NOS: 1 and 3], a polynucleotide of the invention encoding aminopeptidase polypeptide may be obtained using standard cloning and screening methods, such as those for cloning and sequencing chromosomal DNA fragments from bacteria using *Staphylococcus aureus* WCUH 29 cells as starting material, followed by obtaining a full length clone. For example, to obtain a polynucleotide sequence of the invention, such as a sequence given in Table 1 [SEQ ID NOS: 1 and 3], typically a library of clones of chromosomal DNA of *Staphylococcus aureus* WCUH 29 in *E. coli* or some other suitable host is probed with a radiolabeled oligonucleotide, preferably a 17-mer or longer, derived from a partial sequence. Clones carrying DNA identical to that of the probe can then be distinguished using stringent conditions. By sequencing the individual clones thus identified with sequencing primers designed from the original sequence it is then possible to extend the sequence in both directions to determine the full gene sequence. Conveniently, such sequencing is performed using denatured double stranded DNA prepared from a plasmid clone. Suitable techniques are described by Maniatis, T., Fritsch, E.F. and Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989). (see in particular Screening By Hybridization 1.90 and Sequencing Denatured Double-Stranded DNA Templates 13.70). Illustrative of the invention, the polynucleotide set out in Table 1 [SEQ ID NO:1] was discovered in a DNA library derived from *Staphylococcus aureus* WCUH 29.

[0048] The DNA sequence set out in Table 1 [SEQ ID NOS:1] contains an open reading frame encoding a protein having about the number of amino acid residues set forth in Table 1 [SEQ ID NO:2] with a deduced molecular weight that can be calculated using amino acid residue molecular weight values well known in the art. The polynucleotide of SEQ ID NO: 1, between nucleotide number 1 through number 1254 encodes the polypeptide of SEQ ID NO:2. The stop codon begins at nucleotide number 1255 of SEQ ID NO: 1.

[0049] Aminopeptidase of the invention is structurally related to other proteins of the M29 aminopeptidase family, as shown by the results of sequencing the DNA encoding aminopeptidase of the deposited strain. The protein exhibits greatest homology to AMPS\_BACSU (SWISS-PROT: locus AMPS\_BACSU, accession P39762) protein among known proteins. Aminopeptidase of Table 1 [SEQ ID NO:2] has about 61%% identity over its entire length and about 74%% similarity over its entire length with the amino acid sequence of AMPS\_BACSU (SWISS-PROT: locus AMPS\_BACSU, accession P39762) polypeptide.

[0050] The invention provides a polynucleotide sequence identical over its entire length to the coding sequence in Table 1 [SEQ ID NO:1]. Also provided by the invention is the coding sequence for the mature polypeptide or a fragment thereof, by itself as well as the coding sequence for the mature polypeptide or a fragment in reading frame with other coding sequence, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence. The polynucleotide may also contain non-coding sequences, including for example, but not limited to non-coding 5' and 3' sequences, such as the transcribed, non-translated sequences, termination signals, ribosome binding sites, sequences that stabilize mRNA, introns, polyadenylation signals, and additional coding sequence which encode additional amino acids. For example, a marker sequence that facilitates purification of the fused polypeptide can be encoded. In certain embodiments of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, *Proc. Natl. Acad. Sci., USA* 86: 821-824 (1989), or an HA tag (Wilson *et al.*, *Cell* 37: 767 (1984). Polynucleotides of the invention also include, but are not limited to, polynucleotides comprising a structural gene and its naturally associated sequences that control gene expression.

[0051] A preferred embodiment of the invention is a polynucleotide of comprising nucleotide 1 to 1254 or 1257 set forth in SEQ ID NO:1 of Table 1 which encode the aminopeptidase polypeptide.

[0052] The invention also includes polynucleotides of the formula set forth in Table 1 (C)[SEQ ID NO: 1] wherein, at the 5' end of the molecule, X is hydrogen, and at the 3' end of the molecule, Y is hydrogen or a metal, R<sub>1</sub> and R<sub>2</sub> is any nucleic acid residue, and n is an integer between 1 and 1000. Any stretch of nucleic acid residues denoted by either R<sub>n</sub> group, where n is greater than 1, may be either a hetercopolymer or a homopolymer, preferably a heteropolymer.

[0053] The term "polynucleotide encoding a polypeptide" as used herein encompasses polynucleotides that include a sequence encoding a polypeptide of the invention, particularly a bacterial polypeptide and more particularly a polypeptide of the *Staphylococcus aureus* aminopeptidase having the amino acid sequence set out in Table 1 [SEQ ID NO:2]. The term also encompasses polynucleotides that include a single continuous region or discontinuous regions encoding the polypeptide (for example, interrupted by integrated phage or an insertion sequence or editing) together with additional regions, that also may contain coding and/or non-coding sequences.

[0054] The invention further relates to variants of the polynucleotides described herein that encode for variants of the polypeptide having the deduced amino acid sequence of Table 1 [SEQ ID NO:2]. Variants that are fragments of the polynucleotides of the invention may be used to synthesize full-length polynucleotides of the invention.

[0055] Further particularly preferred embodiments are polynucleotides encoding aminopeptidase variants, that have the amino acid sequence of aminopeptidase polypeptide of Table 1 [SEQ ID NO:2] in which several, a few, 5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues are substituted, deleted or added, in any combination. Especially preferred

among these are silent substitutions, additions and deletions, that do not alter the properties and activities of aminopeptidase.

**[0056]** Further preferred embodiments of the invention are polynucleotides that are at least 70% identical over their entire length to a polynucleotide encoding aminopeptidase polypeptide having an amino acid sequence set out in Table 1 [SEQ ID NOS:2 and 4], and polynucleotides that are complementary to such polynucleotides. Alternatively, most highly preferred are polynucleotides that comprise a region that is at least 80% identical over its entire length to a polynucleotide encoding aminopeptidase polypeptide of the deposited strain and polynucleotides complementary thereto. In this regard, polynucleotides at least 90% identical over their entire length to the same are particularly preferred, and among these particularly preferred polynucleotides, those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred among those with at least 95%, and among these those with at least 98% and at least 99% are particularly highly preferred, with at least 99% being the more preferred.

**[0057]** Preferred embodiments are polynucleotides that encode polypeptides that retain substantially the same biological function or activity as the mature polypeptide encoded by the DNA of Table 1 [SEQ ID NO:1].

**[0058]** The invention further relates to polynucleotides that hybridize to the herein above-described sequences. In this regard, the invention especially relates to polynucleotides that hybridize under stringent conditions to the herein above-described polynucleotides. As herein used, the terms "stringent conditions" and "stringent hybridization conditions" mean hybridization will occur only if there is at least 95% and preferably at least 97% identity between the sequences. An example of stringent hybridization conditions is overnight incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ml denatured, sheared salmon sperm DNA, followed by washing the hybridization support in 0.1x SSC at about 65°C. Hybridization and wash conditions are well known and exemplified in Sambrook, *et al.*, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, N.Y., (1989), particularly Chapter 11 therein.

**[0059]** The invention also provides a polynucleotide consisting essentially of a polynucleotide sequence obtainable by screening an appropriate library containing the complete gene for a polynucleotide sequence set forth in SEQ ID NO:1 or SEQ ID NO:3 under stringent hybridization conditions with a probe having the sequence of said polynucleotide sequence set forth in SEQ ID NO: 1 or a fragment thereof; and isolating said DNA sequence. Fragments useful for obtaining such a polynucleotide include, for example, probes and primers described elsewhere herein.

**[0060]** As discussed additionally herein regarding polynucleotide assays of the invention, for instance, polynucleotides of the invention as discussed above, may be used as a hybridization probe for RNA, cDNA and genomic DNA to isolate full-length cDNAs and genomic clones encoding aminopeptidase and to isolate cDNA and genomic clones of other genes that have a high sequence similarity to the aminopeptidase gene. Such probes generally will comprise at least 15 bases. Preferably, such probes will have at least 30 bases and may have at least 50 bases. Particularly preferred probes will have at least 30 bases and will have 50 bases or less.

**[0061]** For example, the coding region of the aminopeptidase gene may be isolated by screening using the DNA sequence provided in SEQ ID NO: 1 to synthesize an oligonucleotide probe. A labeled oligonucleotide having a sequence complementary to that of a gene of the invention is then used to screen a library of cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes to.

**[0062]** The polynucleotides and polypeptides of the invention may be employed, for example, as research reagents and materials for discovery of treatments of and diagnostics for disease, particularly human disease, as further discussed herein relating to polynucleotide assays.

**[0063]** Polynucleotides of the invention that are oligonucleotides derived from the sequences of SEQ ID NOS:1 and/or 2 may be used in the processes herein as described, but preferably for PCR, to determine whether or not the polynucleotides identified herein in whole or in part are transcribed in bacteria in infected tissue. It is recognized that such sequences will also have utility in diagnosis of the stage of infection and type of infection the pathogen has attained.

**[0064]** The invention also provides polynucleotides that may encode a polypeptide that is the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature polypeptide (when the mature form has more than one polypeptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, may allow protein transport, may lengthen or shorten protein half-life or may facilitate manipulation of a protein for assay or production, among other things. As generally is the case *in vivo*, the additional amino acids may be processed away from the mature protein by cellular enzymes.

**[0065]** A precursor protein, having the mature form of the polypeptide fused to one or more prosequences may be an inactive form of the polypeptide. When prosequences are removed such inactive precursors generally are activated. Some or all of the prosequences may be removed before activation. Generally, such precursors are called preproteins.

**[0066]** In sum, a polynucleotide of the invention may encode a mature protein, a mature protein plus a leader sequence (which may be referred to as a preprotein), a precursor of a mature protein having one or more prosequences that are not the leader sequences of a preprotein, or a preproprotein, which is a precursor to a proprotein, having a leader sequence and one or more prosequences, which generally are removed during processing steps that produce

active and mature forms of the polypeptide.

#### Vectors, host cells, expression

[0067] The invention also relates to vectors that comprise a polynucleotide or polynucleotides of the invention, host cells that are genetically engineered with vectors of the invention and the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the invention.

[0068] For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof or polynucleotides of the invention. Introduction of a polynucleotide into the host cell can be effected by methods described in many standard laboratory manuals, such as Davis et al., *BASIC METHODS IN MOLECULAR BIOLOGY*, (1986) and Sambrook et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989), such as, calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction and infection.

[0069] Representative examples of appropriate hosts include bacterial cells, such as streptococci, staphylococci, enterococci *E. coli*, streptomyces and *Bacillus subtilis* cells; fungal cells, such as yeast cells and *Aspergillus* cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, 293 and Bowes melanoma cells; and plant cells.

[0070] A great variety of expression systems can be used to produce the polypeptides of the invention. Such vectors include, among others, chromosomal, episomal and virus-derived vectors, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression system constructs may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides and/or to express a polypeptide in a host may be used for expression in this regard. The appropriate DNA sequence may be inserted into the expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, (*supra*).

[0071] For secretion of the translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the expressed polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

[0072] Polypeptides of the invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography, and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding protein may be employed to regenerate active conformation when the polypeptide is denatured during isolation and or purification.

#### Diagnostic Assays

[0073] This invention is also related to the use of the aminopeptidase polynucleotides of the invention for use as diagnostic reagents. Detection of aminopeptidase in a eukaryote, particularly a mammal, and especially a human, will provide a diagnostic method for diagnosis of a disease. Eukaryotes (herein also "individual(s)"), particularly mammals, and especially humans, infected with an organism comprising the aminopeptidase gene may be detected at the nucleic acid level by a variety of techniques.

[0074] Nucleic acids for diagnosis may be obtained from an infected individual's cells and tissues, such as bone, blood, muscle, cartilage, and skin. Genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification technique prior to analysis. RNA or cDNA may also be used in the same ways. Using amplification, characterization of the species and strain of prokaryote present in an individual, may be made by an analysis of the genotype of the prokaryote gene. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the genotype of a reference sequence. Point mutations can be identified by hybridizing amplified DNA to labeled aminopeptidase polynucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in the electrophoretic mobility of the DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing. See, e.g., Myers et al., *Science*, 230: 1242 (1985). Sequence changes at specific locations also may be revealed by nuclease protection assays, such as RNase and S1

protection or a chemical cleavage method. See, e.g., Cotton et al., *Proc. Natl. Acad. Sci., USA*, 85: 4397-4401 (1988).  
 [0075] Cells carrying mutations or polymorphisms in the gene of the invention may also be detected at the DNA level by a variety of techniques, to allow for serotyping, for example. For example, RT-PCR can be used to detect mutations. It is particularly preferred to use RT-PCR in conjunction with automated detection systems, such as, for example, GeneScan. RNA or cDNA may also be used for the same purpose, PCR or RT-PCR. As an example, PCR primers complementary to a nucleic acid encoding aminopeptidase can be used to identify and analyze mutations. Examples of representative primers are shown below in Table 2.

Table 2

## Primers for amplification of aminopeptidase polynucleotides

<u>SEQ ID NO</u>	<u>PRIMER SEQUENCE</u>
5	5'-CAACAATACG CTGAACTATT AG-3'
6	5'-GAACATCAAT TTAAATTGAT CG-3'

[0076] The invention further provides these primers with 1, 2, 3 or 4 nucleotides removed from the 5' and/or the 3' end. These primers may be used for, among other things, amplifying aminopeptidase DNA isolated from a sample derived from an individual. The primers may be used to amplify the gene isolated from an infected individual such that the gene may then be subject to various techniques for elucidation of the DNA sequence. In this way, mutations in the DNA sequence may be detected and used to diagnose infection and to serotype and/or classify the infectious agent.

[0077] The invention further provides a process for diagnosing, disease, preferably bacterial infections, more preferably infections by *Staphylococcus aureus*, and most preferably disease, such as, infections of the upper respiratory tract (e.g., otitis media, bacterial tracheitis, acute epiglottitis, thyroiditis), lower respiratory (e.g., empyema, lung abscess), cardiac (e.g., infective endocarditis), gastrointestinal (e.g., secretory diarrhoea, splenic abscess, retroperitoneal abscess), CNS (e.g., cerebral abscess), eye (e.g., blepharitis, conjunctivitis, keratitis, endophthalmitis, preseptal and orbital cellulitis, dacryocystitis), kidney and urinary tract (e.g., epididymitis, intrarenal and perinephric abscess, toxic shock syndrome), skin (e.g., impetigo, folliculitis, cutaneous abscesses, cellulitis, wound infection, bacterial myositis) bone and joint (e.g., septic arthritis, osteomyelitis), comprising determining from a sample derived from an individual a increased level of expression of polynucleotide having the sequence of Table 1 [SEQ ID NO: 1]. Increased or decreased expression of aminopeptidase polynucleotide can be measured using any one of the methods well known in the art for the quantitation of polynucleotides, such as, for example, amplification, PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods.

[0078] In addition, a diagnostic assay in accordance with the invention for detecting over-expression of aminopeptidase protein compared to normal control tissue samples may be used to detect the presence of an infection, for example. Assay techniques that can be used to determine levels of a aminopeptidase protein, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

## Antibodies

[0079] The polypeptides of the invention or variants thereof, or cells expressing them can be used as an immunogen to produce antibodies immunospecific for such polypeptides. "Antibodies" as used herein includes monoclonal and polyclonal antibodies, chimeric, single chain, simianized antibodies and humanized antibodies, as well as Fab fragments, including the products of an Fab immunoglobulin expression library.

[0080] Antibodies generated against the polypeptides of the invention can be obtained by administering the polypeptides or epitope-bearing fragments, analogues or cells to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique known in the art that provides antibodies produced by continuous cell line cultures can be used. Examples include various techniques, such as those in Kohler, G. and Milstein, C., *Nature* 256: 495-497 (1975); Kozbor et al., *Immunology Today* 4: 72 (1983); Cole et al., pp. 77-96 in *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc. (1985).

[0081] Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can be adapted to pro-

duce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms such as other mammals, may be used to express humanized antibodies.

[0082] Alternatively phage display technology may be utilized to select antibody genes with binding activities towards the polypeptide either from repertoires of PCR amplified v-genes of lymphocytes from humans screened for possessing anti-aminopeptidase or from naive libraries (McCafferty, J. et al., (1990), *Nature* 348, 552-554; Marks, J. et al., (1992) *Biotechnology* 10, 779-783). The affinity of these antibodies can also be improved by chain shuffling (Clackson, T. et al., (1991) *Nature* 352, 624-628).

[0083] If two antigen binding domains are present each domain may be directed against a different epitope - termed 'bispecific' antibodies.

[0084] The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptides to purify the polypeptides by affinity chromatography.

[0085] Thus, among others, antibodies against aminopeptidase- polypeptide may be employed to treat infections, particularly bacterial infections and especially disease, such as, infections of the upper respiratory tract (e.g., otitis media, bacterial tracheitis, acute epiglottitis, thyroiditis), lower respiratory (e.g., empyema, lung abscess), cardiac (e.g., infective endocarditis), gastrointestinal (e.g., secretory diarrhoea, splenic abscess, retroperitoneal abscess), CNS (e.g., cerebral abscess), eye (e.g., blepharitis, conjunctivitis, keratitis, endophthalmitis, preseptal and orbital cellulitis, dacryocystitis), kidney and urinary tract (e.g., epididymitis, intrarenal and perinephric abscess, toxic shock syndrome), skin (e.g., impetigo, folliculitis, cutaneous abscesses, cellulitis, wound infection, bacterial myositis) bone and joint (e.g., septic arthritis, osteomyelitis).

[0086] Polypeptide variants include antigenically, epitopically or immunologically equivalent variants that form a particular aspect of this invention. The term "antigenically equivalent derivative" as used herein encompasses a polypeptide or its equivalent which will be specifically recognized by certain antibodies which, when raised to the protein or polypeptide according to the invention, interfere with the immediate physical interaction between pathogen and mammalian host. The term "immunologically equivalent derivative" as used herein encompasses a peptide or its equivalent which when used in a suitable formulation to raise antibodies in a vertebrate, the antibodies act to interfere with the immediate physical interaction between pathogen and mammalian host.

[0087] The polypeptide, such as an antigenically or immunologically equivalent derivative or a fusion protein thereof is used as an antigen to immunize a mouse or other animal such as a rat or chicken. The fusion protein may provide stability to the polypeptide. The antigen may be associated, for example by conjugation, with an immunogenic carrier protein for example bovine serum albumin (BSA) or keyhole limpet haemocyanin (KLH). Alternatively a multiple antigenic peptide comprising multiple copies of the protein or polypeptide, or an antigenically or immunologically equivalent polypeptide thereof may be sufficiently antigenic to improve immunogenicity so as to obviate the use of a carrier.

[0088] Preferably, the antibody or variant thereof is modified to make it less immunogenic in the individual. For example, if the individual is human the antibody may most preferably be "humanized"; where the complementarity determining region(s) of the hybridoma-derived antibody has been transplanted into a human monoclonal antibody, for example as described in Jones, P. et al. (1986), *Nature* 321, 522-525 or Tempest et al., (1991) *Biotechnology* 9, 266-273.

[0089] The use of a polynucleotide of the invention in genetic immunization will preferably employ a suitable delivery method such as direct injection of plasmid DNA into muscles (Wolff et al., *Hum Mol Genet* 1992, 1:363, Manthorpe et al., *Hum. Gene Ther.* 1993, 4, 419), delivery of DNA complexed with specific protein carriers (Wu et al., *J Biol Chem.* 1989: 264,16985), coprecipitation of DNA with calcium phosphate (Benvenisty & Reshef, *PNAS USA*, 1986:83,9551), encapsulation of DNA in various forms of liposomes (Kaneda et al., *Science* 1989:243,375), particle bombardment (Tang et al., *Nature* 1992, 356:152, Eisenbraun et al., *DNA Cell Biol* 1993, 12:791) and *in vivo* infection using cloned retroviral vectors (Seeger et al., *PNAS USA* 1984:81,5849).

#### Antagonists and agonists - assays and molecules

[0090] Polypeptides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics. See, e.g., Coligan et al., *Current Protocols in Immunology* 1(2): Chapter 5 (1991).

[0091] The invention also provides a method of screening compounds to identify those which enhance (agonist) or block (antagonist) the action of aminopeptidase polypeptides or polynucleotides, particularly those compounds that are bacteriostatic and/or bacteriocidal. The method of screening may involve high-throughput techniques. For example, to screen for agonists or antagonists, a synthetic reaction mix, a cellular compartment, such as a membrane, cell envelope or cell wall, or a preparation of any thereof, comprising aminopeptidase polypeptide and a labeled substrate or ligand of such polypeptide is incubated in the absence or the presence of a candidate molecule that may be an aminopeptidase agonist or antagonist. The ability of the candidate molecule to agonize or antagonize the aminopeptidase polypeptide is reflected in decreased binding of the labeled ligand or decreased production of product from such sub-

strate. Molecules that bind gratuitously, *i.e.*, without inducing the effects of aminopeptidase polypeptide are most likely to be good antagonists. Molecules that bind well and increase the rate of product production from substrate are agonists. Detection of the rate or level of production of product from substrate may be enhanced by using a reporter system. Reporter systems that may be useful in this regard include but are not limited to colorimetric labeled substrate converted into product, a reporter gene that is responsive to changes in aminopeptidase polynucleotide or polypeptide activity, and binding assays known in the art.

[0092] Another example of an assay for aminopeptidase antagonists is a competitive assay that combines aminopeptidase and a potential antagonist with aminopeptidase-binding molecules, recombinant aminopeptidase binding molecules, natural substrates or ligands, or substrate or ligand mimetics, under appropriate conditions for a competitive inhibition assay. Aminopeptidase can be labeled, such as by radioactivity or a colorimetric compound, such that the number of aminopeptidase molecules bound to a binding molecule or converted to product can be determined accurately to assess the effectiveness of the potential antagonist.

[0093] Potential antagonists include small organic molecules, peptides, polypeptides and antibodies that bind to a polynucleotide or polypeptide of the invention and thereby inhibit or extinguish its activity. Potential antagonists also may be small organic molecules, a peptide, a polypeptide such as a closely related protein or antibody that binds the same sites on a binding molecule, such as a binding molecule, without inducing aminopeptidase-induced activities, thereby preventing the action of aminopeptidase by excluding aminopeptidase from binding.

[0094] Potential antagonists include a small molecule that binds to and occupies the binding site of the polypeptide thereby preventing binding to cellular binding molecules, such that normal biological activity is prevented. Examples of small molecules include but are not limited to small organic molecules, peptides or peptide-like molecules. Other potential antagonists include antisense molecules (see Okano, *J. Neurochem.* 56: 560 (1991); *OLIGODEOXYNUCLEOTIDES AS ANTISENSE INHIBITORS OF GENE EXPRESSION*, CRC Press, Boca Raton, FL (1988), for a description of these molecules). Preferred potential antagonists include compounds related to and variants of aminopeptidase.

[0095] Each of the DNA sequences provided herein may be used in the discovery and development of antibacterial compounds. The encoded protein, upon expression, can be used as a target for the screening of antibacterial drugs. Additionally, the DNA sequences encoding the amino terminal regions of the encoded protein or Shine-Delgarno or other translation facilitating sequences of the respective mRNA can be used to construct antisense sequences to control the expression of the coding sequence of interest.

[0096] The invention also provides the use of the polypeptide, polynucleotide or inhibitor of the invention to interfere with the initial physical interaction between a pathogen and mammalian host responsible for sequelae of infection. In particular the molecules of the invention may be used: in the prevention of adhesion of bacteria, in particular gram positive bacteria, to mammalian extracellular matrix proteins on in-dwelling devices or to extracellular matrix proteins in wounds; to block aminopeptidase protein-mediated mammalian cell invasion by, for example, initiating phosphorylation of mammalian tyrosine kinases (Rosenshine *et al.*, *Infect. Immun.* 60:2211 (1992); to block bacterial adhesion between mammalian extracellular matrix proteins and bacterial aminopeptidase proteins that mediate tissue damage and; to block the normal progression of pathogenesis in infections initiated other than by the implantation of in-dwelling devices or by other surgical techniques.

[0097] The antagonists and agonists of the invention may be employed, for instance, to inhibit and treat disease, such as, infections of the upper respiratory tract (e.g., otitis media, bacterial tracheitis, acute epiglottitis, thyroiditis), lower respiratory (e.g., empyema, lung abscess), cardiac (e.g., infective endocarditis), gastrointestinal (e.g., secretory diarrhoea, splenic abscess, retroperitoneal abscess), CNS (e.g., cerebral abscess), eye (e.g., blepharitis, conjunctivitis, keratitis, endophthalmitis, preseptal and orbital cellulitis, dacryocystitis), kidney and urinary tract (e.g., epididymitis, intrarenal and perinephric abscess, toxic shock syndrome), skin (e.g., impetigo, folliculitis, cutaneous abscesses, cellulitis, wound infection, bacterial myositis) bone and joint (e.g., septic arthritis, osteomyelitis).

#### Vaccines

[0098] Another aspect of the invention relates to a method for inducing an immunological response in an individual, particularly a mammal which comprises inoculating the individual with aminopeptidase, or a fragment or variant thereof, adequate to produce antibody and/ or T cell immune response to protect said individual from infection, particularly bacterial infection and most particularly *Staphylococcus aureus* infection. Also provided are methods whereby such immunological response slows bacterial replication. Yet another aspect of the invention relates to a method of inducing immunological response in an individual which comprises delivering to such individual a nucleic acid vector to direct expression of aminopeptidase, or a fragment or a variant thereof, for expressing aminopeptidase, or a fragment or a variant thereof *in vivo* in order to induce an immunological response, such as, to produce antibody and/ or T cell immune response, including, for example, cytokine-producing T cells or cytotoxic T cells, to protect said individual from disease, whether that disease is already established within the individual or not. One way of administering the gene is by accelerating it into the desired cells as a coating on particles or otherwise. Such nucleic acid vector may comprise DNA,

RNA, a modified nucleic acid, or a DNA/RNA hybrid.

[0099] A further aspect of the invention relates to an immunological composition which, when introduced into an individual capable or having induced within it an immunological response, induces an immunological response in such individual to a aminopeptidase or protein coded therefrom, wherein the composition comprises a recombinant aminopeptidase or protein coded therefrom comprising DNA which codes for and expresses an antigen of said aminopeptidase or protein coded therefrom. The immunological response may be used therapeutically or prophylactically and may take the form of antibody immunity or cellular immunity such as that arising from CTL or CD4+ T cells.

[0100] A aminopeptidase polypeptide or a fragment thereof may be fused with co-protein which may not by itself produce antibodies, but is capable of stabilizing the first protein and producing a fused protein which will have immunogenic and protective properties. Thus fused recombinant protein, preferably further comprises an antigenic co-protein, such as lipoprotein D from *Hemophilus influenzae*, Glutathione-S-transferase (GST) or beta-galactosidase, relatively large co-proteins which solubilize the protein and facilitate production and purification thereof. Moreover, the co-protein may act as an adjuvant in the sense of providing a generalized stimulation of the immune system. The co-protein may be attached to either the amino or carboxy terminus of the first protein.

[0101] Provided by this invention are compositions, particularly vaccine compositions, and methods comprising the polypeptides or polynucleotides of the invention and immunostimulatory DNA sequences, such as those described in Sato, Y. *et al.* Science 273: 352 (1996).

[0102] Also, provided by this invention are methods using the described polynucleotide or particular fragments thereof which have been shown to encode non-variable regions of bacterial cell surface proteins in DNA constructs used in such genetic immunization experiments in animal models of infection with *Staphylococcus aureus* will be particularly useful for identifying protein epitopes able to provoke a prophylactic or therapeutic immune response. It is believed that this approach will allow for the subsequent preparation of monoclonal antibodies of particular value from the requisite organ of the animal successfully resisting or clearing infection for the development of prophylactic agents or therapeutic treatments of bacterial infection, particularly *Staphylococcus aureus* infection, in mammals, particularly humans.

[0103] The polypeptide may be used as an antigen for vaccination of a host to produce specific antibodies which protect against invasion of bacteria, for example by blocking adherence of bacteria to damaged tissue. Examples of tissue damage include wounds in skin or connective tissue caused, e.g., by mechanical, chemical or thermal damage or by implantation of indwelling devices, or wounds in the mucous membranes, such as the mouth, mammary glands, urethra or vagina.

[0104] The invention also includes a vaccine formulation which comprises an immunogenic recombinant protein of the invention together with a suitable carrier. Since the protein may be broken down in the stomach, it is preferably administered parenterally, including, for example, administration that is subcutaneous, intramuscular, intravenous, or intradermal. Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the bodily fluid, preferably the blood, of the individual; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

[0105] While the invention has been described with reference to certain aminopeptidase protein, it is to be understood that this covers fragments of the naturally occurring protein and similar proteins with additions, deletions or substitutions which do not substantially affect the immunogenic properties of the recombinant protein.

#### Compositions, kits and administration

[0106] The invention also relates to compositions comprising the polynucleotide or the polypeptides discussed above or their agonists or antagonists. The polypeptides of the invention may be employed in combination with a non-sterile or sterile carrier or carriers for use with cells, tissues or organisms, such as a pharmaceutical carrier suitable for administration to a subject. Such compositions comprise, for instance, a media additive or a therapeutically effective amount of a polypeptide of the invention and a pharmaceutically acceptable carrier or excipient. Such carriers may include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol and combinations thereof. The formulation should suit the mode of administration. The invention further relates to diagnostic and pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

[0107] Polypeptides and other compounds of the invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

[0108] The pharmaceutical compositions may be administered in any effective, convenient manner including, for instance, administration by topical, oral, anal, vaginal, intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal or intradermal routes among others.

[0109] In therapy or as a prophylactic, the active agent may be administered to an individual as an injectable composition, for example as a sterile aqueous dispersion, preferably isotonic.

[0110] Alternatively the composition may be formulated for topical application for example in the form of ointments, creams, lotions, eye ointments, eye drops, ear drops, mouthwash, impregnated dressings and sutures and aerosols, and may contain appropriate conventional additives, including, for example, preservatives, solvents to assist drug penetration, and emollients in ointments and creams. Such topical formulations may also contain compatible conventional carriers, for example cream or ointment bases, and ethanol or oleyl alcohol for lotions. Such carriers may constitute from about 1% to about 98% by weight of the formulation; more usually they will constitute up to about 80% by weight of the formulation.

[0111] For administration to mammals, and particularly humans, it is expected that the daily dosage level of the active agent will be from 0.01 mg/kg to 10 mg/kg, typically around 1 mg/kg. The physician in any event will determine the actual dosage which will be most suitable for an individual and will vary with the age, weight and response of the particular individual. The above dosages are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

[0112] In-dwelling devices include surgical implants, prosthetic devices and catheters, i.e., devices that are introduced to the body of an individual and remain in position for an extended time. Such devices include, for example, artificial joints, heart valves, pacemakers, vascular grafts, vascular catheters, cerebrospinal fluid shunts, urinary catheters, continuous ambulatory peritoneal dialysis (CAPD) catheters.

[0113] The composition of the invention may be administered by injection to achieve a systemic effect against relevant bacteria shortly before insertion of an in-dwelling device. Treatment may be continued after surgery during the in-body time of the device. In addition, the composition could also be used to broaden perioperative cover for any surgical technique to prevent bacterial wound infections, especially *Staphylococcus aureus* wound infections.

[0114] Many orthopaedic surgeons consider that humans with prosthetic joints should be considered for antibiotic prophylaxis before dental treatment that could produce a bacteremia. Late deep infection is a serious complication sometimes leading to loss of the prosthetic joint and is accompanied by significant morbidity and mortality. It may therefore be possible to extend the use of the active agent as a replacement for prophylactic antibiotics in this situation.

[0115] In addition to the therapy described above, the compositions of this invention may be used generally as a wound treatment agent to prevent adhesion of bacteria to matrix proteins exposed in wound tissue and for prophylactic use in dental treatment as an alternative to, or in conjunction with, antibiotic prophylaxis.

[0116] Alternatively, the composition of the invention may be used to bathe an indwelling device immediately before insertion. The active agent will preferably be present at a concentration of 1µg/ml to 10mg/ml for bathing of wounds or indwelling devices.

[0117] A vaccine composition is conveniently in injectable form. Conventional adjuvants may be employed to enhance the immune response. A suitable unit dose for vaccination is 0.5-5 microgram/kg of antigen, and such dose is preferably administered 1-3 times and with an interval of 1-3 weeks. With the indicated dose range, no adverse toxicological effects will be observed with the compounds of the invention which would preclude their administration to suitable individuals.

[0118] Each reference disclosed herein is incorporated by reference herein in its entirety. Any patent application to which this application claims priority is also incorporated by reference herein in its entirety.

## EXAMPLES

[0119] The examples below are carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. The examples are illustrative, but do not limit the invention.

### Example 1 Strain selection, Library Production and Sequencing

[0120] The polynucleotide having the DNA sequence given in SEQ ID NO:1 was obtained from a library of clones of chromosomal DNA of *Staphylococcus aureus* in *E. coli*. The sequencing data from two or more clones containing overlapping *Staphylococcus aureus* DNAs was used to construct the contiguous DNA sequence in SEQ ID NO:1. Libraries may be prepared by routine methods, for example:

Methods 1 and 2 below.

[0121] Total cellular DNA is isolated from *Staphylococcus aureus* WCUH 29 according to standard procedures and size-fractionated by either of two methods.

**Method 1**

[0122] Total cellular DNA is mechanically sheared by passage through a needle in order to size-fractionate according to standard procedures. DNA fragments of up to 11 kbp in size are rendered blunt by treatment with exonuclease and DNA polymerase, and EcoRI linkers added. Fragments are ligated into the vector Lambda ZapII that has been cut with EcoRI, the library packaged by standard procedures and *E. coli* infected with the packaged library. The library is amplified by standard procedures.

**Method 2**

[0123] Total cellular DNA is partially hydrolyzed with a one or a combination of restriction enzymes appropriate to generate a series of fragments for cloning into library vectors (e.g., RsaI, PstI, AluI, BshI235I), and such fragments are size-fractionated according to standard procedures. EcoRI linkers are ligated to the DNA and the fragments then ligated into the vector Lambda ZapII that have been cut with EcoRI, the library packaged by standard procedures, and *E. coli* infected with the packaged library. The library is amplified by standard procedures.

**Example 2      aminopeptidase Characterization****The determination of expression during infection of a gene from *Staphylococcus aureus***

[0124] Necrotic fatty tissue from a 72hour groin infection or an excised kidney from an 8 day chronic kidney infection of *Staphylococcus aureus* WCUH29 in the mouse is efficiently disrupted and processed in the presence of chaotropic agents and RNAase inhibitor to provide a mixture of animal and bacterial RNA. The optimal conditions for disruption and processing to give stable preparations and high yields of bacterial RNA are followed by the use of hybridisation to a radiolabelled oligonucleotide specific to *Staphylococcus aureus* 16S RNA on Northern blots. The RNAase free, DNAase free, DNA and protein free preparations of RNA obtained are suitable for Reverse Transcription PCR (RT-PCR) using unique primer pairs designed from the sequence of each gene of *Staphylococcus aureus* WCUH29.

**a) Isolation of tissue infected with *Staphylococcus aureus* WCUH29 from a mouse animal model of infection (groin)**

[0125] 10 ml. volumes of sterile nutrient broth (No.2 Oxoid) are seeded with isolated, individual colonies of *Staphylococcus aureus* WCUH29 from an agar culture plate. The cultures are incubated aerobically (static culture) at 37°C for 16-20 hours. 4 week old mice (female, 18g-22g, strain MF1) are each infected by subcutaneous injection of 0.5ml. of this broth culture of *Staphylococcus aureus* WCUH29 (diluted in broth to approximately 10<sup>8</sup> cfu/ml.) into the anterior, right lower quadrant (groin area). Mice should be monitored regularly during the first 24 hours after infection, then daily until termination of study. Animals with signs of systemic infection, i.e. lethargy, ruffled appearance, isolation from group, should be monitored closely and if signs progress to moribundancy, the animal should be culled immediately.

[0126] Visible external signs of lesion development will be seen 24-48h after infection. Examination of the abdomen of the animal will show the raised outline of the abscess beneath the skin. The localised lesion should remain in the right lower quadrant, but may occasionally spread to the left lower quadrant, and superiorly to the thorax. On occasions, the abscess may rupture through the overlying skin layers. In such cases the affected animal should be culled immediately and the tissues sampled if possible. Failure to cull the animal may result in the necrotic skin tissue overlying the abscess being sloughed off, exposing the abdominal muscle wall.

[0127] Approximately 96 hours after infection, animals are killed using carbon dioxide asphyxiation. To minimise delay between death and tissue processing /storage, mice should be killed individually rather than in groups. The dead animal is placed onto its back and the fur swabbed liberally with 70% alcohol. An initial incision using scissors is made through the skin of the abdominal left lower quadrant, travelling superiorly up to, then across the thorax. The incision is completed by cutting inferiorly to the abdominal lower right quadrant. Care should be taken not to penetrate the abdominal wall. Holding the skin flap with forceps, the skin is gently pulled away from the abdomen. The exposed abscess, which covers the peritoneal wall but generally does not penetrate the muscle sheet completely, is excised, taking care not to puncture the viscera.

[0128] The abscess/muscle sheet and other infected tissue may require cutting in sections, prior to flash-freezing in liquid nitrogen, thereby allowing easier storage in plastic collecting vials.

**b) Isolation of tissue infected with *Staphylococcus aureus* WCUH29 from a murine model of hematogenous pyelonephritis.**

[0129] Overnight cultures of *S. aureus* WCUH29 were started from single colonies in 5 ml of tryptic soy broth (TSB) and grown at 37°C with shaking. The cultures were then washed twice in sterile phosphate-buffered saline (PBS) and diluted to an A600 = 0.3. Male CD-1 mice (18 - 20g) were infected with 0.2 ml of this suspension by tail vein inoculation using a 30g needle attached to a tuberculin syringe. Each mouse receives approximately  $4 \times 10^7$  bacteria in this fashion. Mice are monitored daily for signs of illness, and usually within 48 hours show signs of lethargy, ruffled fur, sluggishness; animals which appear moribund are euthanized prior to the end of the experiment.

[0130] All animals are euthanized via carbon dioxide overdose seven days post-infection. The animal is placed on its back and swabbed with ethanol, and then with RNAZap, and instruments are swabbed as well. The abdominal cavity is opened and the kidneys aseptically removed, cut into four pieces, and placed in cryovials which are immediately frozen in liquid nitrogen.

**c) Isolation of *Staphylococcus aureus* WCUH29 RNA from infected tissue samples**

[0131] 4-6 infected tissue samples (each approx 0.5-0.7g) in 2ml screw-cap tubes are removed from -80°C storage into a dry ice ethanol bath. In a microbiological safety cabinet the samples are disrupted individually whilst the remaining samples are kept cold in the dry ice ethanol bath. To disrupt the bacteria within the tissue sample 1ml of TRIzol Reagent (Gibco BRL, Life Technologies) is added followed by enough 0.1mm zirconia/silica beads to almost fill the tube, the lid is replaced taking care not to get any beads into the screw thread so as to ensure a good seal and eliminate aerosol generation. The sample is then homogenised in a Mini-BeadBeater Type BX-4 (Biospec Products). Necrotic fatty tissue is strain treated for 100 seconds at 5000 rpm in order to achieve bacterial lysis. *In vivo* grown bacteria require longer treatment than *in vitro* grown *Staphylococcus aureus* which are disrupted by a 30 second bead-beat.

[0132] After bead-beating the tubes are chilled on ice before opening in a fume-hood as heat generated during disruption may degrade the TRIzol and release cyanide.

[0133] 200 microlitres of chloroform is then added and the tubes shaken by hand for 15 seconds to ensure complete mixing. After 2-3 minutes at room temperature the tubes are spun down at 12,000 x g, 4°C for 1 minutes and RNA extraction is then continued according to the method given by the manufacturers of TRIzol Reagent i.e.: The aqueous phase, approx 0.6 ml, is transferred to a sterile eppendorf tube and 0.5 ml of isopropanol is added. After 10 minutes at room temperature the samples are spun at 12,000 x g, 4°C for 10 minutes. The supernatant is removed and discarded then the RNA pellet is washed with 1 ml 75% ethanol. A brief vortex is used to mix the sample before centrifuging at 7,500 x g, 4°C for 5 minutes. The ethanol is removed and the RNA pellet dried under vacuum for no more than 5 minutes. Samples are then resuspended by repeated pipetting in 100 microlitres of DEPC treated water, followed by 5-10 minutes at 55°C. Finally, after at least 1 minute on ice, 200 units of Rnasin (Promega) is added.

[0134] RNA preparations are stored at -80°C for up to one month. For longer term storage the RNA precipitate can be stored at the wash stage of the protocol in 75% ethanol for at least one year at -20°C.

[0135] Quality of the RNA isolated is assessed by running samples on 1% agarose gels. 1 x TBE gels stained with ethidium bromide are used to visualise total RNA yields. To demonstrate the isolation of bacterial RNA from the infected tissue 1 x MOPS, 2.2M formaldehyde gels are run and vacuum blotted to Hybond-N (Amersham). The blot is then hybridised with a <sup>32</sup>P labelled oligonucleotide probe specific to 16s rRNA of *Staphylococcus aureus* (K. Greisen, M. Loeffelholz, A. Purohit and D. Leong. J.Clin. (1994) Microbiol. 32 335-351). An oligonucleotide of the sequence: 5'-gctcctaaaagggtactccaccggc-3' is used as a probe. The size of the hybridising band is compared to that of control RNA isolated from *in vitro* grown *Staphylococcus aureus* WCUH29 in the Northern blot. Correct sized bacterial 16s rRNA bands can be detected in total RNA samples which show extensive degradation of the mammalian RNA when visualised on TBE gels.

**d) The removal of DNA from *Staphylococcus aureus* WCUH29-derived RNA**

[0136] DNA is removed from 73 microlitre samples of RNA by a 15 minute treatment on ice with 3 units of DNAase I, amplification grade (Gibco BRL, Life Technologies) in the buffer supplied with the addition of 200 units of Rnasin (Promega) in a final volume of 90 microlitres.

[0137] The DNAase is inactivated and removed by treatment with TRIzol LS Reagent (Gibco BRL, Life Technologies) according to the manufacturers protocol. DNAase treated RNA is resuspended in 73 microlitres of DEPC treated water with the addition of Rnasin as described in Method 1.

**e) The preparation of cDNA from RNA samples derived from infected tissue**

**[0138]** 10 microlitre samples of DNAase treated RNA are reverse transcribed using a SuperScript Preamplification System for First Strand cDNA Synthesis kit (Gibco BRL, Life Technologies) according to the manufacturers instructions. 1 nanogram of random hexamers is used to prime each reaction. Controls without the addition of SuperScriptII reverse transcriptase are also run. Both +/-RT samples are treated with RNaseH before proceeding to the PCR reaction

**f) The use of PCR to determine the presence of a bacterial cDNA species**

**[0139]** PCR reactions are set up on ice in 0.2ml tubes by adding the following components: 45 microlitres PCR SUPERMIX (Gibco BRL, Life Technologies); 1 microlitre 50mM MgCl<sub>2</sub>, to adjust final concentration to 2.5mM; 1 microlitre PCR primers (optimally 18-25 basepairs in length and designed to possess similar annealing temperatures), each primer at 10mM initial concentration; and 2 microlitres cDNA.

**[0140]** PCR reactions are run on a Perkin Elmer GeneAmp PCR System 9600 as follows: 5 minutes at 95 °C, then 50 cycles of 30 seconds each at 94 °C, 42 °C and 72 °C followed by 3 minutes at 72 °C and then a hold temperature of 4 °C. (the number of cycles is optimally 30-50 to determine the appearance or lack of a PCR product and optimally 8-30 cycles if an estimation of the starting quantity of cDNA from the RT reaction is to be made); 10 microlitre aliquots are then run out on 1% 1 x TBE gels stained with ethidium bromide with PCR product, if present, sizes estimated by comparison to a 100 bp DNA Ladder (Gibco BRL, Life Technologies). Alternatively if the PCR products are conveniently labelled by the use of a labelled PCR primer (e.g. labelled at the 5' end with a dye) a suitable aliquot of the PCR product is run out on a polyacrylamide sequencing gel and its presence and quantity detected using a suitable gel scanning system (e.g. ABI Prism™ 377 Sequencer using GeneScan™ software as supplied by Perkin Elmer).

**[0141]** RT/PCR controls may include +/- reverse transcriptase reactions, 16s rRNA primers or DNA specific primer pairs designed to produce PCR products from non-transcribed *Staphylococcus aureus* WCUH29 genomic sequences.

**[0142]** To test the efficiency of the primer pairs they are used in DNA PCR with *Staphylococcus aureus* WCUH29 total DNA. PCR reactions are set up and run as described above using approx. 1 microgram of DNA in place of the cDNA and 35 cycles of PCR.

**[0143]** Primer pairs which fail to give the predicted sized product in either DNA PCR or RT/PCR are PCR failures and as such are uninformative. Of those which give the correct size product with DNA PCR two classes are distinguished in RT/PCR: 1. Genes which are not transcribed *in vivo* reproducibly fail to give a product in RT/PCR; and 2. Genes which are transcribed *in vivo* reproducibly give the correct size product in RT/PCR and show a stronger signal in the +RT samples than the signal (if at all present) in -RT controls

Annex to the description

[0144]

5

## SEQUENCE LISTING

## (1) GENERAL INFORMATION

10

(i) APPLICANT: SmithKline Beecham Corporation & SmithKline  
Beecham Plc

15

(ii) TITLE OF THE INVENTION: NOVEL aminopeptidase

20

(iii) NUMBER OF SEQUENCES: 7

## (iv) CORRESPONDENCE ADDRESS:

25

(A) ADDRESSEE: SmithKline Beecham, Corporate Intellectual  
Property

(B) STREET: Two New Horizons Court

(C) CITY: Brentford

(D) STATE: Middlesex

30

(E) COUNTRY: United Kingdom

(F) ZIP: TW8 9EP

## (v) COMPUTER READABLE FORM:

35

(A) MEDIUM TYPE: Diskette

(B) COMPUTER: IBM Compatible

(C) OPERATING SYSTEM: DOS

(D) SOFTWARE: FastSEQ for Windows Version 2.0

40

## (vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:

(B) FILING DATE:

45

(C) CLASSIFICATION:

## (vii) PRIOR APPLICATION DATA:

50

(A) APPLICATION NUMBER:

(B) FILING DATE:

55

## (viii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: CONNELL, Anthony Christopher  
 (B) REGISTRATION NUMBER: 5630 & 26758  
 (C) REFERENCE/DOCKET NUMBER: GM10055

## (ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: +44 1279 644 395  
 (B) TELEFAX: +44 181 975 6294  
 (C) TELEX:

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1257 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GTGGGAATTT	TGACTAATTA	CAAAGAAAAG	TTACAACAAT	ACGCTGAACT	ATTAGTTAAG	60
GTTGGTATGA	ATGTTCAACC	AAAACAGCCA	GTATTTATTA	GATCATCAGT	AGAAATATTA	120
GAATTAACGC	ATTTAATTGT	AGAAGAAGCA	TATCATTGTG	GCGCTTCAGA	TGTACGGGTT	180
GTATATAGCG	ATCCAACCTCT	TAAAACGCTT	AAATTTGAAA	ATGAATCTGT	GGAACATTTT	240
GCTAATCATG	AGATAAAATC	ATATGATGTT	GAAGCGAGAA	TGGATTACGT	TAAACGTGGC	300
GCTGCTAATT	TAGCACTTAT	TAGTGAAGAT	CCTGATTTAA	TGGATAGTAT	CGATAGCCAA	360
AAATTACAAG	CATTTCAACA	ACAAAACGCA	CGCGCATTTA	AAGGATATAT	GGAAAAGTGTA	420
CAAAAAAATC	AATTTCCATG	GGTCGTAGCT	GCTTTTCCAT	CTAAAGCTTG	GGCGAAGAGA	480
GTATATCCTG	AACTATCTGT	AGAAGAAGCA	TATATTAAGT	TTATTGACGA	GGTATTTGAT	540
ATTGTGCGAA	TTGATGGTAA	CGATCCAGTT	GAAAATTGGC	GTCAACACAT	CGCTAATTTA	600
AGTGTTTACG	CTCAATATTT	ACAACAAAAA	AATTACCATG	CGTTGCATGA	TGTATCTGAA	660
GGTACTGATT	TAACAGTATG	TTTAGCGAAA	AATCATATCT	GGGAAGATGC	TACAAGTTAT	720
GTTAACGGCG	AGGAACAAGC	TTTTATTGCT	AATATACCAA	CTGAAGAAGT	ATTTACAGCA	780
CCAGATCGAA	ATAGAGTACA	TGGATATGTC	ACTAATAAAT	TGCCTTTAAG	TTATAATGGA	840
ACGATTATCG	ATCAATTTAA	ATTGATGTTT	AAAGATGGAG	AAATTATTGA	TTTTTCAGCT	900
GAAAAAGGTG	AAGCGGTATT	GAAAGATTTA	ATCAATACCG	ATGAAGGTTT	AAGAAGATTA	960
GGTGAAGTAG	CGTTAGTACC	TGATGATTCA	CCGATATCAA	ATCGTAATAC	CATTTTTTAC	1020
AATACTTTAT	TTGATGAAAA	TGCAGCATGT	CATTTAGCGA	TTGGATCTGC	TTACGCCCTT	1080

AATATTCAAG GTGGAACGGA AATGACTGTT GAGGAAAAGA TTGCAAGCGG ATTAAATGAT 1140  
 TCAAACGTAC ATGTCGATTT CATGATTGGT AGTAGTGATT TGACTATTTA TGGCATATTC 1200  
 5 GAAGATGGTT CAAAAGAACT AGTATTTGAA AATGGAAATT GGGCATCAAC ATTTTAA 1257

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 418 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Val Gly Ile Leu Thr Asn Tyr Lys Glu Lys Leu Gln Gln Tyr Ala Glu  
 1 5 10 15  
 25 Leu Leu Val Lys Val Gly Met Asn Val Gln Pro Lys Gln Pro Val Phe  
 20 25 30  
 Ile Arg Ser Ser Val Glu Ile Leu Glu Leu Thr His Leu Ile Val Glu  
 35 40 45  
 30 Glu Ala Tyr His Cys Gly Ala Ser Asp Val Arg Val Val Tyr Ser Asp  
 50 55 60  
 Pro Thr Leu Lys Thr Leu Lys Phe Glu Asn Glu Ser Val Glu His Phe  
 65 70 75 80  
 35 Ala Asn His Glu Ile Lys Ser Tyr Asp Val Glu Ala Arg Met Asp Tyr  
 85 90 95  
 Val Lys Arg Gly Ala Ala Asn Leu Ala Leu Ile Ser Glu Asp Pro Asp  
 100 105 110  
 40 Leu Met Asp Ser Ile Asp Ser Gln Lys Leu Gln Ala Phe Gln Gln Gln  
 115 120 125  
 Asn Ala Arg Ala Phe Lys Gly Tyr Met Glu Ser Val Gln Lys Asn Gln  
 130 135 140  
 45 Phe Pro Trp Val Val Ala Ala Phe Pro Ser Lys Ala Trp Ala Lys Arg  
 145 150 155 160  
 Val Tyr Pro Glu Leu Ser Val Glu Glu Ala Tyr Ile Lys Phe Ile Asp  
 165 170 175  
 50 Glu Val Phe Asp Ile Val Arg Ile Asp Gly Asn Asp Pro Val Glu Asn  
 180 185 190  
 Trp Arg Gln His Ile Ala Asn Leu Ser Val Tyr Ala Gln Tyr Leu Gln

55

	195		200		205
	Gln Lys Asn Tyr His Ala Leu His Asp Val Ser Glu Gly Thr Asp Leu				
5	210		215		220
	Thr Val Cys Leu Ala Lys Asn His Ile Trp Glu Asp Ala Thr Ser Tyr				
	225		230		235
	Val Asn Gly Glu Glu Gln Ala Phe Ile Ala Asn Ile Pro Thr Glu Glu				
10		245		250	255
	Val Phe Thr Ala Pro Asp Arg Asn Arg Val His Gly Tyr Val Thr Asn				
	260		265		270
15	Lys Leu Pro Leu Ser Tyr Asn Gly Thr Ile Ile Asp Gln Phe Lys Leu				
	275		280		285
	Met Phe Lys Asp Gly Glu Ile Ile Asp Phe Ser Ala Glu Lys Gly Glu				
	290		295		300
20	Ala Val Leu Lys Asp Leu Ile Asn Thr Asp Glu Gly Ser Arg Arg Leu				
	305		310		315
	Gly Glu Val Ala Leu Val Pro Asp Asp Ser Pro Ile Ser Asn Arg Asn				
		325		330	335
25	Thr Ile Phe Tyr Asn Thr Leu Phe Asp Glu Asn Ala Ala Cys His Leu				
	340		345		350
	Ala Ile Gly Ser Ala Tyr Ala Phe Asn Ile Gln Gly Gly Thr Glu Met				
	355		360		365
30	Thr Val Glu Glu Lys Ile Ala Ser Gly Leu Asn Asp Ser Asn Val His				
	370		375		380
	Val Asp Phe Met Ile Gly Ser Ser Asp Leu Thr Ile Tyr Gly Ile Phe				
	385		390		395
35	Glu Asp Gly Ser Lys Glu Leu Val Phe Glu Asn Gly Asn Trp Ala Ser				
		405		410	415
	Thr Phe				

40

## (2) INFORMATION FOR SEQ ID NO:3:

45

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 858 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

50

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

55

5 CTAATTANCA AAGAAAAGTT ACAACAATAC GCTGAACTAT TAGTTAAGGT TGGTATGAAT 60  
 GTTCAACCAA AACAGCCAGT ATTTATTAGA TCATCAGTAG AAACATTAGA ATTAACGCAT 120  
 TTAATTGTAG AAGAAGCATA TCATTGTGGC GCTTCAGATG TACGTGTTGT ATATAGCGAT 180  
 CCAACCTTAA AACGCTTAAA ATTTGAAAAT GAATCTGTGG AACATTTTGC TAATCATGAG 240  
 10 ATAAAATCAT ATGATGTTGA AGCGAGAATG GATTACGTTA AACGTGGCGC TGCTAATTTA 300  
 GCACTTATTA GTGAAGATCC TGATTTAATG GATGGTATCG ATAGCCAAAA ATTACAAGCA 360  
 TTTCAACAAC AAAACGCACG CGCATTTAAA GGATATATGG AAAGTGTACA AAAAAATCAA 420  
 TTTCCATGGG TCGTAGCTGC TTTTCCATCT AAAGCTTGGG CGAAGAGAGT ATATCCTGAA 480  
 15 CTATCTGTAG AAGAAGCATA TATTAAGTTT ATTGACGAGG TATTTGATAT TGTGCGAATT 540  
 GATGGTAACG ATCCAGTTGA AAATTGGCGT CAACACATCG CTAATTTAAG TGTTTACGCT 600  
 CAATATTTAC AACAAAAAAA TTACCATGCG TTGCATGATG TATCTGAAGG TACTGATTTA 660  
 ACAGTATGTT TAGCGAAAAA TCATATCTGG GAAGATGCTA CAAGTTATGT TAACGGCGAG 720  
 20 GAACAAGCTT TTATTGCTAA TATACCAACT GAAGAAGTAT TTACAGCACC AGATCGAAAT 780  
 AGAGTACATG GATATGTCAC TAATAAATTG CCTTTAAATT TATTTGGAGC GATATTCGAT 840  
 CAATTTAAAT TGATGTTC 858

25 (2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- 30 (A) LENGTH: 285 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

40 Ile Xaa Lys Glu Lys Leu Gln Gln Tyr Ala Glu Leu Leu Val Lys Val  
 1 5 10 15  
 Gly Met Asn Val Gln Pro Lys Gln Pro Val Phe Ile Arg Ser Ser Val  
 20 25 30  
 45 Glu Thr Leu Glu Leu Thr His Leu Ile Val Glu Glu Ala Tyr His Cys  
 35 40 45  
 Gly Ala Ser Asp Val Arg Val Val Tyr Ser Asp Pro Thr Leu Lys Arg  
 50 55 60  
 50 Leu Lys Phe Glu Asn Glu Ser Val Glu His Phe Ala Asn His Glu Ile  
 65 70 75 80  
 Lys Ser Tyr Asp Val Glu Ala Arg Met Asp Tyr Val Lys Arg Gly Ala  
 85 90 95

55

	Ala	Asn	Leu	Ala	Leu	Ile	Ser	Glu	Asp	Pro	Asp	Leu	Met	Asp	Gly	Ile
				100					105					110		
5	Asp	Ser	Gln	Lys	Leu	Gln	Ala	Phe	Gln	Gln	Gln	Asn	Ala	Arg	Ala	Phe
			115					120					125			
	Lys	Gly	Tyr	Met	Glu	Ser	Val	Gln	Lys	Asn	Gln	Phe	Pro	Trp	Val	Val
		130					135					140				
10	Ala	Ala	Phe	Pro	Ser	Lys	Ala	Trp	Ala	Lys	Arg	Val	Tyr	Pro	Glu	Leu
	145					150					155					160
	Ser	Val	Glu	Glu	Ala	Tyr	Ile	Lys	Phe	Ile	Asp	Glu	Val	Phe	Asp	Ile
					165				170						175	
15	Val	Arg	Ile	Asp	Gly	Asn	Asp	Pro	Val	Glu	Asn	Trp	Arg	Gln	His	Ile
				180					185					190		
	Ala	Asn	Leu	Ser	Val	Tyr	Ala	Gln	Tyr	Leu	Gln	Gln	Lys	Asn	Tyr	His
		195						200					205			
20	Ala	Leu	His	Asp	Val	Ser	Glu	Gly	Thr	Asp	Leu	Thr	Val	Cys	Leu	Ala
		210					215					220				
	Lys	Asn	His	Ile	Trp	Glu	Asp	Ala	Thr	Ser	Tyr	Val	Asn	Gly	Glu	Glu
25	225					230					235					240
	Gln	Ala	Phe	Ile	Ala	Asn	Ile	Pro	Thr	Glu	Glu	Val	Phe	Thr	Ala	Pro
					245					250					255	
	Asp	Arg	Asn	Arg	Val	His	Gly	Tyr	Val	Thr	Asn	Lys	Leu	Pro	Leu	Asn
30				260					265					270		
	Leu	Phe	Gly	Ala	Ile	Phe	Asp	Gln	Phe	Lys	Leu	Met	Phe			
			275					280					285			

35 (2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

40 (A) LENGTH: 22 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

50 CAACAATACG CTGAACTATT AG

22

(2) INFORMATION FOR SEQ ID NO: 6:

55

(2) INFORMATION FOR SEQ ID NO: 6:

55

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

GAACATCAAT TTAAATTGAT CG

22

## (2) INFORMATION FOR SEQ ID NO:7:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GCTCCTAAAA GGTTACTCCA CCGGC

25

## Claims

1. An isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of:

- (a) a polynucleotide having at least a 70% identity to a polynucleotide encoding a polypeptide comprising the amino acid sequence of SEQ ID NO:2;
- (b) a polynucleotide having at least a 70% identity to a polynucleotide encoding the same mature polypeptide expressed by the aminopeptidase gene contained in the *Staphylococcus aureus* of the deposited strain;
- (c) a polynucleotide encoding a polypeptide comprising an amino acid sequence which is at least 70% identical to the amino acid sequence of SEQ ID NO:2;
- (d) a polynucleotide which is complementary to the polynucleotide of (a), (b) or (c); and
- (e) a polynucleotide comprising at least 15 sequential bases of the polynucleotide of (a), (b), (c) or (d).

2. The polynucleotide of Claim 1 wherein the polynucleotide is DNA.

3. The polynucleotide of Claim 1 wherein the polynucleotide is RNA.

4. The polynucleotide of Claim 2 comprising the nucleic acid sequence set forth in SEQ ID NO:1.

5. The polynucleotide of Claim 2 comprising nucleotide 1 to 1252 set forth in SEQ ID NO:1.

6. The polynucleotide of Claim 2 which encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:2.

7. A vector comprising the polynucleotide of Claim 1.

8. A host cell comprising the vector of Claim 7.

5 9. A process for producing a polypeptide comprising: expressing from the host cell of Claim 8 a polypeptide encoded by said DNA.

10 10. A process for producing an aminopeptidase polypeptide or fragment comprising culturing a host of claim 8 under conditions sufficient for the production of said polypeptide or fragment.

11. A polypeptide comprising an amino acid sequence which is at least 70% identical to the amino acid sequence of SEQ ID NO:2.

12. A polypeptide comprising an amino acid sequence as set forth in SEQ ID NO:2.

13. An antibody against the polypeptide of claim 11.

14. An antagonist which inhibits the activity or expression of the polypeptide of claim 11.

20 15. A method for the treatment of an individual in need of aminopeptidase polypeptide comprising: administering to the individual a therapeutically effective amount of the polypeptide of claim 11.

25 16. A method for the treatment of an individual having need to inhibit aminopeptidase polypeptide comprising: administering to the individual a therapeutically effective amount of the antagonist of Claim 14.

17. A process for diagnosing a disease related to expression or activity of the polypeptide of claim 11 in an individual comprising:

- 30 (a) determining a nucleic acid sequence encoding said polypeptide, and/or  
(b) analyzing for the presence or amount of said polypeptide in a sample derived from the individual.

18. A method for identifying compounds which interact with and inhibit or activate an activity of the polypeptide of claim 11 comprising:

35 contacting a composition comprising the polypeptide with the compound to be screened under conditions to permit interaction between the compound and the polypeptide to assess the interaction of a compound, such interaction being associated with a second component capable of providing a detectable signal in response to the interaction of the polypeptide with the compound;  
40 and determining whether the compound interacts with and activates or inhibits an activity of the polypeptide by detecting the presence or absence of a signal generated from the interaction of the compound with the polypeptide.

45 19. A method for inducing an immunological response in a mammal which comprises inoculating the mammal with aminopeptidase polypeptide of claim 11, or a fragment or variant thereof, adequate to produce antibody and/or T cell immune response to protect said animal from disease.

50 20. A method of inducing immunological response in a mammal which comprises delivering a nucleic acid vector to direct expression of aminopeptidase polypeptide of claim 11, or fragment or a variant thereof, for expressing said aminopeptidase polypeptide, or a fragment or a variant thereof *in vivo* in order to induce an immunological response to produce antibody and/ or T cell immune response to protect said animal from disease.